

# Cloud Seeding Program

## 2017 Operations Report

July 2018

## Introduction

Snowy Hydro Limited (Snowy Hydro) relies on precipitation falling over the catchments of the Snowy Mountains to supply water for the production of hydroelectricity. Cloud seeding over this area is used to enhance snowfall, ultimately leading to increased runoff and amount of water available to produce energy.

Cloud seeding operations undertaken by Snowy Hydro are authorised by the *Snowy Mountains Cloud Seeding Act 2004* (NSW) (SMCS Act).

The SMCS Act mandates that cloud seeding operations may only be carried out in accordance with an Environmental Management Plan (EMP) approved by the Minister administering the *Environmental Planning and Assessment Act 1979*, and the Minister administering Part 4 of the *National Parks and Wildlife Act 1974* (the 'relevant Ministers').

The Cloud Seeding Program EMP was formally approved by the relevant Ministers on 5 July 2013 following consultation with experts from the Office of Environment and Heritage (OEH), NSW Environment Protection Authority (EPA) and National Parks and Wildlife Service (NPWS).

The SMCS Act requires Snowy Hydro, by 31 March in each year, to report on cloud seeding operations during the previous year to the relevant Ministers and to the EPA. The report must include details of compliance with the EMP and details of research monitoring the impact of seeding agents on the environment.

The EPA is appointed to review each report on cloud seeding operations, and report the findings of the review and any resulting recommendations to the Board of the EPA and the relevant Ministers.

Snowy Hydro submitted the Cloud Seeding Program 2017 Annual Compliance Report to the relevant Ministers and the EPA in March 2018. The report demonstrated that Snowy Hydro has carried out cloud seeding operations in accordance with the SMCS Act and has complied with all obligations set out within the EMP, with the exception of potable water sampling at one site which not not completed due to safety reasons. Importantly, the 2017 Annual Compliance Report confirmed there continues to be no evidence of any significant adverse environmental impacts associated with cloud seeding activities.

The subsequent EPA review<sup>1</sup> published in June 2018 supported these findings.

The key points of the 2017 Annual Compliance Report that are described fully in the following sections of this report:

- Operations, including the duration over which cloud seeding occurred and the total amount of cloud seeding agent released over the season;
- Meteorological monitoring, including controls to ensure precipitation falls as snow to at least 1400 metres during cloud seeding operations and assessment of downwind impacts; and
- Environmental monitoring, including summary statistics of the monitoring program and details of research monitoring the impact of seeding agents on the environment.

Finally, the findings and recommendations of the EPA review are provided.

---

<sup>1</sup> Report on the Findings of the NSW Environment Protection Authority's Review of the Snowy Hydro Limited Cloud Seeding Program: 2017 Annual Compliance Report. Available from <https://www.epa.nsw.gov.au/legislation/snowy-hydro-cloud-seed.htm>

---

## Operations

### Target area

The SMCS Act states that the area primarily targeted for increased precipitation must be land within the Snowy water catchment.

An area of approximately 2110 km<sup>2</sup> was targeted during 2017 cloud seeding operations. Figure 1 shows both the target area and the Snowy water catchment boundaries.

### Hours of operation

The SMCS Act stipulates that operations are only to be carried out when precipitation is likely to fall as snow to at least 1400 metres. Consequently, cloud seeding operations take place throughout the cool-season months, typically between May and September.

In 2017, a total of 95 hours and 48 minutes of cloud seeding occurred between 8 July 2017 and 14 September 2017.

### Seeding agent and method of discharge

Silver iodide is the approved seeding agent. It is used as the ice nucleating material because it has similar physical properties to an ice crystal. It is also insoluble in water and non-bioavailable. In 2017, approximately 36.4 kg of silver iodide was dispersed over the 2110 km<sup>2</sup> target area.

Land-based aerosol generators are the approved method to disperse the seeding material. The seeding agent is released by up to 23 ground-based generators located along the western perimeter of the target area when suitable atmospheric conditions are present<sup>2</sup>.

### Operational incidents

There were no accidents or breakdowns resulting in spillage of cloud seeding agents, fuel, or failure of controls specified in the EMP.

---

<sup>2</sup> For explanation of how cloud seeding works and the atmospheric conditions required for cloud seeding operations, see <http://www.snowyhydro.com.au/our-energy/cloud-seeding/>.




Figure 1: Map showing the Snowy water catchment (the area approved for cloud seeding) and the area which is primarily targeted for increased precipitation from cloud seeding operations (approximately 2110 km<sup>2</sup>)

## Meteorological Monitoring

### Snow level criterion

The SMCS Act mandates that the discharge of the seeding agent is to be carried out at a time when increased precipitation is likely to fall as snow to at least 1400 metres above sea level.

Weather balloons are released at three hourly intervals before and during cloud seeding operations to monitor atmospheric conditions. Cloud seeding does not commence, or is suspended, if the freezing level measured over the catchment is higher than 1600 metres. This is to ensure precipitation falls as snow to at least 1400 metres.

Additional controls are implemented if the freezing level is between 1550 and 1600 metres, including monitoring live camera feeds and verifying conditions with personnel within the target area.

During 2017:

- Cloud seeding operations did not commence when the freezing level over the catchment was greater than 1600 metres.
- There was one instance where the freezing level rose above 1600 metres during operations at which stage the event was suspended.
- Freezing levels during operations were observed to be between 1550-1600 metres during one seeding event, triggering additional controls:
  - Cameras within the target area were monitored and no rain was observed falling at locations between 1327 and 1481 metres.
  - External personnel within the target area at an elevation of approximately 1350 metres confirmed snow precipitation.

### Downwind impact

The cloud seeding program has been designed so that additional precipitation from cloud seeding falls over the target area. A component of the design is routine monitoring of precipitation to identify any possible effects of cloud seeding extending outside the target area.

Data from Bureau of Meteorology and Snowy Hydro weather stations provide the basis for comparison of the temporal and spatial variability of precipitation across the region during the winter months, both before and after cloud seeding operations commenced in 2004.

Analyses of precipitation amounts over 1990-2017 continue to show no evidence of an effect from cloud seeding on precipitation downwind of the target area.<sup>3</sup> This supports results of previous, independent analyses by the Natural Resources Commission (NRC).

---

<sup>3</sup> NRC Review of SPERP Annual Report 2011 (July 2012), available from [www.nrc.nsw.gov.au/cloud-seeding](http://www.nrc.nsw.gov.au/cloud-seeding).

## Environmental Monitoring

### Environmental chemistry

Snowy Hydro has monitored silver concentrations in a range of environmental matrices at potential accumulation zones within and around the target area since the commencement of cloud seeding over the Snowy Mountains in 2004.

The objectives of the monitoring program are to detect significant increases in the concentrations of silver compared with baseline concentrations, and to assess concentrations of silver compared with agreed guideline values of 0.1 mg/L for potable water and 1 mg/kg for all other matrices.

The EMP prescribes the number of sampling sites for each matrix and area, the replicates collected and analysed for each site and the sampling frequency. Once environmental samples are collected, they are sent to a laboratory for chemical analysis. The results are independently audited and analysed statistically.

Analyses of silver concentrations from samples collected prior to the commencement of cloud seeding in 2004 through to 2017 continues to show no evidence that cloud seeding has contributed to increased levels of silver in any of the areas, or in any of the environmental matrices monitored.

In accordance with the EMP, potable water was the only matrix sampled following the cessation of the 2017 season. The EMP lists twelve sites for potable water sampling. Eleven of the twelve sites were sampled following the 2017 cloud seeding season; despite best endeavours, safe access could not be obtained for the final site which is on private property. The total number of samples analysed in 2017 along with summary statistics of silver concentrations is shown in Table 1 for potable water samples. All measurements are well below the relevant guideline values.

**Table 1: Summary of silver concentrations in 2017 potable water samples (ng/L). The guideline value for silver in potable water is 100,000 ng/L**

<i>Matrix</i>	<i>Number of samples</i>	<i>Minimum</i>	<i>Mean</i>	<i>Maximum</i>	<i>Guideline value</i>
Potable Water	33	0.46	1.56	7.98	100,000

### Aquatic ecology

Analyses of data collected following the 2013 season showed no evidence of any difference over time in the impairment of the macroinvertebrate assemblages or multivariate structure of edge or riffle assemblages which could be related to cloud seeding. Aquatic macroinvertebrates sampling will therefore take place after the 2018 cloud seeding season, in accordance with the EMP.

### Environmental fate study

Researchers from the University of Queensland were enlisted by SHL in 2012 to investigate the environmental fate of silver and indium<sup>4</sup> in the Snowy Water Catchment deposited from the atmosphere,

<sup>4</sup> Indium was considered as part of this study although the approved, inert, tracing agent indium sesquioxide has not been released since 2011

---

either from long-range sources unrelated to cloud seeding, or released during cloud seeding operations. This investigation was concluded in 2014 and no evidence was found that cloud seeding had increased silver concentrations in the environment. The results of this study have now been published in international peer reviewed literature.

Silver is naturally present in the atmosphere, soil and sediments of the Snowy Mountains. This originates from natural weathering of rocks and from dust sourced from metal-containing landscapes upwind of the Snowy Mountains. In addition, metals released hundreds of kilometres away from industrial sources such as fossil fuel combustion and metal production are transported through the atmosphere and deposited in the Snowy Mountains. This process has occurred for at least the past 100 years.

The University of Queensland research team provided a briefing on the investigation findings to OEH and EPA in April 2015. SHL considered these recommendations, and developed a report addressing proposed changes to the Cloud Seeding EMP. The recommendations do not impact the annual potable water sampling campaign. SHL presented the proposed changes to the EPA in 2017 as part of the required 5-yearly review of the EMP. The EPA, on behalf of the Minister for the Environment and the Minister for Planning, completed the review of the EMP in July 2018. SHL will implement any subsequent adaptations to the environmental monitoring program in advance of the major sampling campaign that will be conducted at the conclusion of the 2018 cloud seeding season.

## EPA Review

Following submission of the 2017 Annual Compliance Report, the EPA published its review in June 2018. The EPA concluded:

- *“Snowy Hydro Limited has complied with all of its obligations as detailed in the Act*
- *Snowy Hydro Limited has complied with all of its obligations as detailed in the Environmental Management Plan for cloud seeding operations approved by the relevant Ministers on 5 July 2013, with the exception of sampling at one potable water sampling site.*
- *Snowy Hydro Limited has complied with all of its obligations as detailed in the Protection of the Environment Operations Act 1997*
- *analysis of potable water sampling carried out during the 2017 cloud seeding season showed similar silver concentrations to those in previous years and remain several orders of magnitude below relevant guideline values”.*

The recommendations were:

- *“future reviews are also undertaken in consultation with input from partner agencies of the National Parks and Wildlife Service and the Office of Environment and Heritage*
- *Snowy Hydro Limited continues to pursue research opportunities on the cloud seeding operations in the Snowy Mountains*
- *the outcomes of this review are communicated to the relevant Ministers.”*

## Conclusion

The 2017 Annual Compliance Report detailing cloud seeding operations and activities through 2017 was submitted to the relevant Ministers and EPA in March 2018. The EPA reviewed the report and confirmed Snowy Hydro has complied with all obligations set out in the SMCS Act and detailed within the EMP through the reporting period. There continues to be no evidence of any significant adverse environmental impacts associated with cloud seeding activities.

For more information on Snowy Hydro’s Cloud Seeding Program please refer to our website, <http://www.snowyhydro.com.au/our-energy/cloud-seeding/>.





